THE PLANT DISEASE REPORTER

Issued By

Division of Mycology and Disease Survey



#### BUREAU OF PLANT INDUSTRY

### UNITED STATES DEPARTMENT OF AGRICULTURE

Supplement 129


Index to Supplement 121 - 128

December 31, 1940

(Issued December 31, 1941)



The Plant Disease Reporter is issued as a service to plant pathologists throughout the United States. It contains reports, summaries, observations, and comments submitted voluntarily by qualified observers. These reports often are in the form of suggestions, queries, and opinions, frequently purely tentative, offered for consideration or discussion rather than as matters of established fact. In accepting and publishing this material the Division of Mycology and Disease Survey serves merely as an informational clearing house. It does not assume responsibility for the subject matter.



# INDEX OF ORGANISMS AND NON-PARASITIC DISEASES

in

# THE PLANT DISEASE REPORTER

SUPPLEMENTS 121 - 128

Prepared by Nellie W. Nance

Plant Disease Reporter Supplement 129

December 31, 1940

Parris, G. K. A check list of fungi, bacteria, nematodes and viruses occurring in Hawaii, and their hosts, Supplement 121:1-91.

Darker, G. D. A brief host index of some plant pathogens and virus diseases in eastern Asia. Supplement 122:93-123.

Cooke, Wm. Bridge. Preliminary host index to fungi of Mount Shasta, California. Supplement 123:125-133.

Barss, Howard P. Proceedings of the third National Plant Nematode Conference. Supplement 124:135-150.

Maneval, W. E. Some recent records of plant pathogens in Missouri. Supplement 125:151-164. (These are listed alphabetically).

Melchers, L. E. and Alvin E. Lowe. The reaction of sorghum varieties and hybrids to milo disease. Supplement 126:165-175.

Edson, H. A., and Jessie I. Wood. Crop losses from plant diseases in the United States in 1939. Supplement 127:177-209.

Nance, Nellie W. Diseases of plants in the United States in 1939. Supplement 128:210-378.

A

Actinomyces sp., sweetpotato, 189, 271.

scabies, garden beet, 253. potato, 191, 287.

Adelopus balsamicola, Abies lasiocarpa, 346.

Albinism, Brassica oleracea var. broccoli, 255. peanut, 325.

Albinism, pepper, 261.

Albugo bliti, Amaranthus retroflexus, 350.

candida, Capsella bursa-pastoris,

horseradish, 285.

radish, 285.

ipomoeae-panduranae, Ipomoea hederacea, 363.

Armillaria mellea, Acer spp., 336. Albugo ipomoeae-panduranae, Ipomoea black oak, 342. pandurata, 364. cherry, 313. sweetpotato, 271. Genistra sp., 361. occidentalis, spinach, 293. mountain ash, 342. platensis, Boerhavia erecta, 353. Norway maple, 342. tragopogonis, salsify, 294. Paeonia spp., 368. Alkali scil, strawberry, 208. peach, 295. Alternaria spp., castor bean, 332. Rosa spp., 372. Cornus asperifolia, 339. sugar maple, 342. cotton, 325, 326. Thuja spp., 349. flax, 228. Ascochyta sp., Clematis spp., 356. garden beet, 253. Erica sp., 359. Hydrangea sp., 364. green peas, 198, 199. Jussiaea decurrens, 365. pea, 284. pepper, 260. white sweet clover, 240. summer squash, 269. caulicola, sweet clover, 240. tobacco, 330. graminicola, barley, 226. tomato, 192, 275. imperfecta, alfalfa, 238. blight, pea, 283. lethalis, sweet clover, 240. brassicae, Brassica oleracea var. pinodella, pea, 284. broccoli, 254. pinodella, vetch, 242. cabbage, 256. pisi, pea, 284, see also Mycosphaecauliflower, 255. rella pinodes. turnip, 259. rhei, rhubarb, 286. carotae, carrot, 270. Asia, eastern, host index of plant cucumerina, cantaloupe, 263. pathogens and virus diseases cucumber, 265. in, 93-123. squash, 267. Aspergillus spp., cotton, 325. watermelon, 262. field corn, 234. dianthi, Dianthus caryophyllus, sweetpotato, 271. 358. niger, field corn, 234 herculea, cabbage, 256. grape, 322. cauliflower, 255. onion, 248. radish, 285. ochraceus, field corn, 234. radicina, carrot, 270. Asperisporium caricae, papaya, 300. solani, eggplant, 286. Aster yellows (virus), Calendula pepper, 260. officinalis, 354. potato, 191, 287. celery, 252. tomato, 194, 274. lettuce, 273. tenuis, Bryophyllum pinnatum, 353. Atropellis piniphila, Pinus ponderosa, tomato, tomato, 194, 274. 348. Anguina tritici, wheat, 230. Angular yellow spot, peach, 299. Aphanomyces cochlioides, sugar beet, 333. Bacillus sp., sugar beet, 333. euteiches, pea, 283. Bacteria, in Hawaii and their hosts, Aphelenchoides fragariae, 76-77. Chrysanthemum sop., 356. Bacterial wilt, tobacco, 202. fragariae, strawberry, 208, 301.

Arachnopeziza aurelia, Quercus sp.,

342.

Bacterium dissolvens, see Erwinia

carotovora.

Bacterium mori, see Phytomonas mori.
Baldwin spot, apple, see bitter pit.
Basidiophora entospora, Callistephus
chinensis, 354.
Big vein (virus), lettuce, 273.
Bitter pit (non-par.), apple, 310.
Black bundle, corn, 187.

Black-end (non-par.), pear, 318.
Black heart (non-par.), celery, 252.
Black leaf spot, Diospyros spp.,

339.

Black ring (virus), Brassica oleracea var. broccoli, 255. Brassica oleracea var. italica, 255.

cauliflower, 256.

Black root and root rot (undet), strawberry, 208, 304.

Black spot (non-par.), garden beet, 253.

Blakemore yellows, see leaf variegation.

Blast (non-par.), oats, 185, 226.

Blight, hops, 328.

Blight (bacterial), Ulmus parvifolia, 345.

Blight (non-par.), garden beet, 253.

Blight (undet.), Matthiola spp., 367.

Blight and dieback (undet.), Thuja spp., 349.

Blossom blight (non-par.), cherry, 315.

Blossom-end rot (non-par.), pepper, 261.

tomato, 193, 195, 278.

Blotchy ripening (non-par.), tomato, 278.

Blue-green algae, blue grass, 242. clover, 242.

Botryosphaeria ribis, red currant, 319.

Botryosporium pulchrum, tomato, 275. Botrytis sp., Antirrhinum majus, 351.

Delphinium spp., 357.

Dianthus caryophyllus, 358.

grape, 322.

Lilium spp., 366.

onion, 248.

Botrytis sp., Paeonia spp., 368. allii, onion, 248.

cinerea, Euphorbia spp., 360.

Gladiolus spp., 362.

lettuce, 272.

Lilium spp., 366.

strawberry, 302.

elliptica, Lilium spp., 366.

tulipae, Tulipa spp., 376.

Boydia insculpta, Ilex spp., 340.

Breaking of blossoms, Dianthus caryophyllus, 359.

Bremia lactucae, lettuce, 272.

Sonchus oleraceus, 374.

Brown rcot rot (undet.), tobacco, 332. Bubakia crotonis, Croton monanthogynus, 357.

Bumpy peach (virus), peach, 298. Bunchy top, eggplant, 286. tomato, 279.

C

Calico (undet.), peach, 299. Canker, Boysenberry, 320.

hops, 328.

Canker-stain; London planetree, 340.

Capnodium sp., Citrus sinensis, 300. Pinus strobus, 348.

Cephalosporium sp., Diospyros spp., 339.

acremonium, corn, 187.

field corn, 234.

Ceratostomella sp., London plane tree, 341.

Platanus occidentalis, 341. fimbriata, sweetpotato, 189, 271.

ulmi, Ulmus spp., 344.

Cercospora sp., Ambrosia trifida, 350.

Catalpa spp., 338.

Hibiscus rosa-sinensis, 363.

Hymenocallis occidentalis, 364.

peanut, 325.

strawberry, 302.

Vernonia baldwinii, 376.

Viburnum spp., 376.

acalyphae, Acalyphya ostryaefolia, 350.

althaeina, Althaea rosea, 350. cotton, 326.

Cercospora antipus, Lonicera sp., 366. Cercosporella rubi, dewberry, 320. apii, celery, 250. Cerebella andropogonis, Axonopus apii carotae, carrot, 270. compressus, 244. Tricholaena repens, 247. apii-pastinacae, parsnip, 280. arborescentis, Hydrangea arbores-Check list of fungi, bacteria, nematodes and viruses occurring in cens, 364. Hawaii, 1-91. asparagi, asparagus, 252. beticola, Beta vulgaris macrorhiza, Chlorosis, Buxus sempervirens, 354. Camellia japonica, 355. garden beet, 253. Catalpa spp., 338. sugar beet, 200, 333. cherry, 315. capsici, pepper, 260. Choisya ternata, 356. Citrus grandis, 300. Citrus sinensis, 300. catalpae, Catalpa spp., 338. chenopodii, Chenopodium hybridum, 355. Gladiolus spp., 262. citrullina, watermelon, 262. huckleberry, 304. cruenta, cowpea, 242. Ilex spp., 340. flagellaris, Phytolacca americana, Italian prune, 313. 370. Ligustrum spp., 366. fusca, pecan, 324. peach, 206. gaurae, Gaura biennis, 361. peanut, 325. medicaginia, alfalfa, 238. Quercus palustris, 343. meliae, Melia azedarach, 367. Rosa spo., 373. minima, pear, 317. sour cherry, 315. moricola, Morus alba, 311. Spiraea spp., 374. nicotianae, tobacco, 330. sweet cherry, 315. oryzae, rice, 228. Thuja spp., 349. pachypus, Helianthus sop., 363. Choanephora sp., Hibiscus spp., 363. personata, peanut, 325. conjuncta, cotton, 326. pittospori, Pittosporum spp., 371. Iponoea spp., 364. rhododendri, Rhododendron okra, 271. pea, 284. catawbiense, 372. ricinella, castor bean, 332. Petunia spp., 370. rosicola, Rosa spp., 372. cucurbitarum, cowpea, 242. rosigena, Rosa spp., 372. okra, 271. smilacina, Smilax bona-nox, 374. Cintractia subinclusa, Carex rostrata, sorghi, Sorghum halepense, 246. 355. varia, Viburnum opulus, 376. Cladosporium, grape, 322. variicolor, Paeonia spp., 369. carpophilum, peach, 206, 295. vignae, cowpea, 242. plum, 316, violae, Viola spp., 377. cucumerinum, cantaloupe, 263. zebrina, Trifolium hybridum, 241. squash, 267. summer squash, 269. Trifolium pratense, 241. Cercosporella albo-maculans, effusum, pecan, 324. fulvum, tomato, 275. Brassica chinensis, 254. petsai cabbage, 259. herbarum, pepper, 260. paeoniae, Paeonia spp., 369. turnip, 259. herpotrichoides, wheat, 230, 231. pisicola, pea, 284. pastinacae, parsnip, 280. Claviceps sp., Agropyron smithii, 243. Agropyron trachycaulum, 243. rubi, blackberry, 319.

Claviceps sp., Elymus spp., 245. Coniothyrium wernsdorffiae, Rosa spp., purpurea, Agropyron repens, 243. 372. Agropyron spicatum, 243. Copper injury, cherry, 315. barley, 226. Copper russetting, apple, 311. rye, 182, 229. Cork (non-par.), apple, 310. wheat, 230. cherry, 315. Clitocybe tabescens, tung-oil tree, Corticium microsclerotia, fig, 301. 325. sasakii, Cynodon dactylon, 244. Coccomyces hiemalis, cherry, 207, 313. solani, bean, 281. lutescens, chokecherry, 335. cabbage, 256. prunophorae, plum, 316. cauliflower, 255. Cold injury, tobacco, 332. Cichorium endivia, 261. Coleosporium solidaginis, Callistephus cotton, 326. chinensis, 354. lettuce, 272. Pinus resinosa, 348. potato, 287. Colletotrichum sp., African succulent sweetpotato, 271. Euphorbias, 360. tomato, 275. Euphorbia fimbriata, 360. stevensii, fig, 301. pepper, 260. Pittosporum spp., 371. antirrhini, Antirrhinum majus, Coryneum beijerinckii, almond, 324. 351. apricot, 312. circinans, onion, 248. peach, 295. destructivum, alfalfa, 238. prune, 312. Melilotus alba, 240. Prunus emarginata, 314. falcatum, sugar cane, 334. sweet cherry, 314. fruit rot, tamato, 276. cardinale, Monterey cypress, 346. Cracked stem (non-par.), celery, 252. graminicolum, Agrostis alba, 243. alfalfa, 238. Cracking (non-par.), cherry, 315. red and white clovers, 241. Crazy top, almond, 324. rye, 183, 229. cotton, 327. Sorghum vulgare sudanense, 246. Crinkle (virus), Pelargonium sop., 369. strawberry, 208, 304. wheat, 231. hibisci, Hibiscus spo., 363. Crinkle leaf, cotton, 327. Cronartium sp., Quercus alba, 336. lagenarium, cantaloupe, 263. cucumber, 266. Quercus macrocarpa, 336. squash, 267. Quercus rubra, 336. cerebrum, Pinus banksiana, 348. watermelon, 262. lindemuthianum, bean, 196, 281. comptoniae, Pinus spp., 348. harknessii, Pinus ponderosa, 348. cowpea, 242. green beans, 196. quercuum, Pinus spp., 348. ribicola, black current, 319. lini, flax, 228. currant, 319. nigrua, pepper, 260. phomoides, tomato, 275. gooseberry, 318. spinaciae, spinach, 293. Ribes, 348, 372. white pine, 348, 372. trifolii, alfalfa, 238. Crown and shoot rot, milo, see milo violae-tricoloris, Viola disease. papilionacea, 377. Crown girdle (undet.), pear, 318. Coniothyrium sp., Ulmus spp., 344. concentricum, Yucca filamentosa, tung-oil tree, 325. Crown rot (undet.), garden beet, 254. 377.

Cryptosporella viticola, grape, 323. Cucumber mosaic (virus), celery, 252. sweet pepper, 261. Cumminsiella sanguinea, Berberis aouifolium, 353. Curly top (virus), bean, 283. Beta vulgaris macrorhiza, 254. cantaloupe, 265. cucumber, 267. dry beans, 197. garden beet, 253. pepper, 261. potato, 291. squash, 268. sugar beet, 200, 334. swiss chard, 254. tomato, 193, 195, 278. Curvularia lunata, rice, 228. Cuscuta sp., Ambrosia trifida, 350. Antirrhinum majus, 351. eggplant, 286. Helianthus spp., 363. tomato, 275. arvensis, alfalfa, 238. Cutting rot, Azalea spp., 353. Cyanospora albicedrae, Juniperus mexicana, 347. Cycloconium oleaginum, olive, 312. Cylindrosporium sp., Amorpha fruticosa, 351. Fraxinus spp., 340. Fraxinus toumeyi, 340. consociatum, Acer glabrum, 336. filipendulae, Spiraea spp., 374. griseum, Sapindus drummondii, 373. montenegrinum, Trollius albiflorus, 376. oculatum, Populus deltoides, 341. spiraeicola, Spiraea lucida, 374. Cytospora sp., Acer spp., 336. Picea spp., 347. Pice pungens, 347. Populus spp., 341. Ulmus spp., 344. chrysosperma, Populus spp., 341. Salix spp., 343. sacchari, sugar cane, 334.

Damping-off, broadleaves, 335.

conifers, 335.

Damping-off, Pinus ponderosa, 335, 346. Pinus resinosa, 346. sugar beet, 200. Dark center (non-par.), turnip, 259. Dasyscypha sp., Pseudotsuga taxifolia, 348. Dead hills, hops, 328. Deficiency diseases, cotton, 201, 327. tobacco, 202. Dendrophoma obscurans, strawberry, 302. Diachea leucopoda, strawberry, 302. Diamond canker (undet.), prune, 313. Diaporthe batatatis, sweetpotato, 271. phaseolorum, lima bean, 280. sojae, soybean, 240. Dibotryon morbosum, cherry, 314. prune, 313. Prunus maritima, 316. Didymella applanata, raspberry, 320. Didymellina macrospora, Iris sp., 364. Dieback, Azalea spp., 353. Boysenberry, 320. Calluna vulgaris, 355. cherry, 315. filbert, 324. prune, 313. Robinia pseudoacacia, 343. Rosa spp., 373. Taxus cuspidata, 349. Diplocarpon earliana, strawberry, 208, 302. rosae, Rosa spp., 373. Diplodia sp., Allium sativum, 250. corn, 186, 187, 233. Rosa spp., 373. sweet corn, 188, 237. caryogena, Carya cordiformis, 338. gossypina, cotton, 325, 326. longispora, Quercus prinus, 342. natalensis, Citrus limonia, 300. tubericola, sweetpotato, 272. zeae, corn, 186, 187. Discosia artocreas, Cercis canadensis, 339. Ditylenchus dipsaci, alfalfa, 238. Allium sativum, 250. narcissus, 140. onion, 248. strawberry, 302. Dormant hills, hops, 328. Dothichiza populea, Populus spp., 341.

Dothiorella sp., tung-oil tree, 325. Erwinia carotovora, cauliflower, 255. ulmi, Ulmus spp., 344. celery, 251. Drought injury, birch, 338. field corn, 235. cabbage, 256. Iris sp., 364. cherry, 315. onion, 248. field corn, 237. potato, 287. horsechestnut, 337. squash, 268. ornamental trees and shrubs, tobacco, 330. 336. tomato, 275. pea, 285. lapsa, field corn, 235. peach, 300. sorghum, 230. pear, 318. phytophthora, potato, 190, 288. Picea spp., 348. tracheiphila, cantaloupe, 264. Pinus sylvestris, 348. cucumber, 266. wheat, 233. watermelon, 262. Drought and heat injury, oats, 226. winter squash, 268. potato, 293. Erysiphe sp., Euphorbia sp., 360. strawberry, 304. cichoracearum, cantaloupe, 264. Drought spot (non-par.), prune, 313. cucumber, 266. Dwarf, strawberry, 208. Helianthus spp., 363. Phlox spp., 370. E salsify, 294. squash, 268. Ear rots, corn, 186, 187, 234. sunflower, 238. Elsinoë ampelina, grape, 323. Taraxacum spp., 375. veneta, blackberry, 319. winter squash, 268. Boysenberry, 320. Zinnia elegans, 377. dewberry, 320. graminis, barley, 181, 226. raspberry, 320. Dactylis glomerata, 245. youngberry, 322. oats, 225. Elytroderma deformans, Pinus rye, 229. ponderosa, 348. turnip, 259. wheat, 231. Endothia parasitica, chestnut, 338. Entomosporium sp., crabapple, 336. polygoni, bean, 281. hawthorn, 336. Brassica oleracea var. napothuemenii, Crataegus spp., 339. brassica, 254. Entyloma sp., Physalis sp., 370. cabbage, 257. dahliae, Dahlia spp., 357. cowpea, 242. nymphaea, Nymphaea odorata, 368. Delphinium spp., 358. green beans, 196. oryzae, rice, 228. Erwinia amylovora, apple, 204, 205, pea, 284. Phaseolus aureus, 280. 304. Polygonum buxiforme, 371. Crataegus spp., 339. red clover, 241. pear, 203, 317. Exobasidium burtii, Azalea spp., 352. Pyracantha sp., 371. vaccinii, Azalea spp., 352. quince, 301. Exosporium concentricum, Euonymus aroideáe, Delphinium spp., 358. japonica, 360. Zantedeschia sp., 377. carotovora, Brussels sprouts, 259. cabbage, 257. Fabraea sp., crabapple, 336. carrot, 270.

celery, 251.

Fabraea sp., hawthorn, 336. Fusarium spp., cotton, 325, 326. maculata, Crataegus spp., 339. Dahlia spb., 357. pear, 203, 317. Dianthus caryophyllus, 358. quince, 301. field corn, 235, 236. False blossom (virus), cranberry, Freesia sp., 361. Gladiolus spp., 362. 322. Fasciation, yew, see witches! green peas, 198, 199. broom. Iris kaempferi, 364. Filiform mosaic (cucurbit virus), Narcissus sp., 367. tomato, 279. oats, 225. Fomes sp., horsechestnut, 337. onion, 248... annosus, Kalmia latifolia, 365. pea, 283, 284. Picea spp., 347. Pelargonium spp., 369. applanatus, Robinia pseudoacacia, pepper, 260. potato, 190, 288. 343. fraxinophilus, Fraxinus lanceolata, Primula sp., 371. 340. sorghum, 230. igniarius, Salix spp., 343. spinach, 293. Foot rots, corn, 186, 187, 236. summer squash, 269. rye, 183: sweet corn, 188, 237. wheat, 178, 179. sweetpotato, 272. Frenching (non-par.), tobacco, 332. Taxus cus pidata, 349. Frost injury, apple, 311. tobacco, 330. cherry, 315. wheat, 231. peach, 300. pear, 318. white sweet clover, 240. winter squash, 269. plum, 317. Zucchini squash, 269. prune, 313. apii, celery, 251. Syringa vulgaris, 375. apii pallidum, celery, 251. Frost killing, Abies concolor, 335. avenaceum, peach, 295. broadleaves, 335. potato, 288. conifers, 335. bulbigenum batatas, sweetpotato, Pseudotsuga taxifolia, 335. 189, 272. Fruit gumming (non-par.), peach, 299. bulbigenum lycopersici, tomato, 192, 194, 275. Fruit pox, tomato, 278. Fruit rots, strawberry, 208. bulbigenum niveum, watermelon, 262. tomato, 192, 194. bulbigenum niveum f.2, cantaloupe, Fumago soo., Ilex spo., 340. Quercus spp., 342. bulbigenum tracheiphilum, cowpea, 242. conglutinans, cabbage, 257. vagans, Fagus spp., 339. cauliflower, 255. Fungi, of Mount Shasta, California, host index to, 125-133. callistephi, Callistephus in Hawaii and their hosts, 37chinensis, 354. heterosporium, Axonopus compressus, Fugarium sp., Agrestis spp., 243. 244. javanicum, summer squash, 269. asparagus, 252. barley, 180. lini, flax, 228. moniliforme, cotton, 325, 326. bean, 281. cantaloupe, 264. field corn, 235.

sugar cane, 334.

Fusarium nivale, Agrostis spp., 243. Gloeodes pomigena, apple, 305. orthoceras longius, strawberry, pear, 317. 302. Gloeosporium spp., Acer spp., 336. orthoceras pisi, pea, 284. Nephthytis sp., 368. oxysporum, sweetpotato, 189, 272. Rosa spp., 373. oxysporum f.1, potato, 288. apocryptum, Acer spp., 336. oxysporum f.2., sweetpotato, 189, catalpae, Catalpa spp., 338. fagi, Fagus spp., 339. oxysporum f.8, pea, 284. perennans, apple, see Neofabraea oxysporum gladioli, Gladiolus perennans. spp., 362. ulmi, Ulmus americana, 344. oxysporum medicaginis, alfalfa, Ulmus pumila, 344. Glomerella cingulata, apple, 204, 205, oxysporum nicotianae, tobacco, 305. 330. Ligustrum spp., 366. perniciosum, mimosa, 337. pepper, 260. radicicola, potato, 288. Symphoricarpos spp., 374. root rot, carrot, 270. gossypii, cotton, 201, 325, 326. solani eumartii, potato, 288. piperata, pepper, 260. solani var. martii f.2, pea, 283. Gnomonia leptostyla, black walnut, 340. solani var. martii f.3, bean, 281. ulmea, Ulmus spp., 344. spinaciae, spinach, 293. veneta, Platanus spp., 341. vasinfectum, cotton, 201, 325, Quercus spp., 342. 326. sycamore, 342. vasinfectum zonatum f.l, onion, Graphiola phoenicis, Phoenix sp., 370. 248. Phoenix canariensis, 370. wilt, casaba melon, 265. Green mosaic (virus), sugar cane, 334. cotton, 146, 147. Greene mottle, apple, 310. Guignardia aesculi, horsechestnut, 337. honeydew melon, 265. Fusicladium eriobotryae, loquat, bidwellii, grape, 209, 323. Parthenocissus quinquefolia, 369. 301. saliciperdum, Salix spp., 343. Parthenocissus tricuspidata, 369. tremulae, Populus spp., 341. vaccinii, cranberry, 322. Fusicoccum sp., Pyracantha sp., 371. Gymnoconia peckiana, black raspberry, Fusisporium rubi, dewberry, 321. 320. raspberry, 321. blackberry, 319. wild blackberry, 319. G Gymnosporangium sp., Amelanchier sp., 350. apple, 204, 205, 306. Ganoderma sp., Colutea arborescens, Crataegus spp., 336, 339. 356. Juniperus monosperma, 347. Gibberella spp., field corn, 236. Malus spp., 336. sweet corn, 237. betheli, Juniperus scopulorum, 347. saubinetii, see G. zeae. zeae, barley, 181, 226. clavariaeforme, Juniperus communis depressa, 347. field corn, 235. oats, 225. clavipes, apple, 305. Crataegus spp., 339.

flowering crab, 340.

Juniperus spp., 347.

rye, 229.

wheat, 178, 179, 231.

Girdle (non-par.), garden beet, 253.

Gymnosporangium clavipes, quince, Helminthosporium turcicum, sorghum, 301. exiguum, Juniperus mexicana, 347. Sorghum vulgare sudanense, 246. floriforme, Juniperus spp., 347. sweet corn, 237. globosum, apple; 306. Heterodera marioni, Althaea rosea, 350. Crataegus spp., 339. American-Egyptian cotton, 143. Juniperus spp., 346, 347. Antirrhinum majus, 351. bean, 138, 147, 281. juniperi-virginianae, apple, 306. juniperi-virginianae, flowering Brabham peas, 139. crab, 340. cantaloupe, 264. Juniperus spp., 346, 347. carrot, 270. Juniperus chinensis, 347. celery, 251. chemical treatment of soil, 150. juniperinum, Juniperus communis depressa, 347. Chinese lily, 1/41. nelsoni, Juniperus spp., 347. cigar-wrapper tobacco, 143. clover, 241. cern, 139. H cotton, 139, 146, 201, 326. Hail injury, pear, 318. cowpea, 138, 139, 242. crop plants, 142. prune, 313. Hair sprout, potato, 291. cucumber, 266. Hawaii, check list of fungi, bacteria, effect of crop rotation, 145. nematodes, and viruses in, eggplant, 286. Euonymus spp., 360. 1-91. Heart rot, sugar beet, 334. Gardenia jasminoides, 361. Heat canker, flax, 228. Gelsemium sempervirens, 361. Hest injury, cabbage, 256. Gladiolus spp., 362. globe artichoke, 269. field corn, 237. pea, 285. grasses, :143. · Tsuga spp., 349. hops, 329. wheat, 233. huckleberry, 304. Helminthosporium spp., Agrostis spp., Ipomoea purpurea, 364. Lespedeza stipulacea, 139. 243. lettuce, 273. Allium sativum, 250. barley, 180. Ligustrum spp., 366. Ligustrum lucidum compactum, 366. field corn, 236. Poa spp., 246. lima bean, 139. Nancy Hall sweet potato, 141. Stenotaphrum secundatum, 246. undetermined Gramineae, 247. Nicotiana spp., 140. whe t, 231. okra, 271. Paeonia spn., 369. avenae, oats, 225. Paper Thite narcissus, 141. bromi, Bromus spp., 244. foot rot, barley 180. pea, .234. gramineum, barley, 180, 227. peach, 296. peanut, 139. oryzae, see Ophiobolus miyabeanus. Petunia spp., 370. papulosum, ample, 306. Pinus lambertiana, 139. sativum, barley, 227. wheat, 231. potate, 142, 145; 288. turcicum, corn, 187. resistance in Nicotiana, 139. field corn, 236. Rosa spp., 373.

Heterodera marioni, snap bean, 139. K soybean, 139. strawberry, 302. Kabatiella caulivora, clover, 241. sweetpotato, 272. Kuehneola uredinis, blackberry, 319. tobacco, 145, 330. tomato, 139, 141, 146, 149, 192, 276. turnip, 259: Leaf blight (non-par.), Arbutus Ulmus spp., 345. menziesii, 337. Viburnum spo., 376. Leaf blotch (sterile fungus), rice, Viola spp., 377. 229. watermelon, 150, 262. Leaf cast, Abies spp., 346. weeds, 142. Leaf-casting yellows (virus), peach, 298. marioni infestation, 148. Leaf curl (undet.), prune, 313. Heterosporium echinulatum, Dianthus Leaf curl (virus), raspberry, 322. caryophyllus, 358. Leaf drop, prune, 313. gracile, Belamcanda chinensis, Robinia pseudoacacia, 343. 353. Leaf mottle, Fagus spp., 339. Iris sp., see Didymellina Leaf necrosis (undet.), turnip, 259. macrospora. Leafroll (virus), potato, 190, 291. phlei, Phleum pratense, 246. Leaf scorch, Acer spp., 337. variabile, spinach, 293. Fagus spp., 340. Hopperburn (leafhopper), potato, 191, prune, 313. 292. Leaf scorch and blight (non-par.), Host index, fungi of Mount Shasta, Pseudotsuga taxifolia, 349. California, 125-133. Leaf spot (non-par.), Camellia japonica, plant pathogens and virus diseases, eastern Asia, Phleum pratense, 246. 92-123. Leaf spotting (undet.), prune, 313. Hyalopsora polypodii, Cystopteris Leaf spotting and defoliation (undet.), peach, 299. sp., 357. Hypodermella nervisequia, Pseudotsuga Le f variegation, strawberry, 208, 303. Leptosphaeria coniothyrium, blackberry, taxifolia, 348. Hypoxylon sp., Platanus occidentalis, 319. Boysenberry, 320. 341. atropunctata, Quercus spp., 342. dewberry, 320. gooseberry, 318. raspberry, 321. Ι Rosa spp., 373. youngberry, 322. Index, host, fungi of Mount Shasta, Leptosphaeria salvinii, rice, 228. California, 125-133. Leptothyrium acerinum, Acer saccharum, plant pathogens and virus diseases, eastern Asia, 336. periclymeni, Lonicera sp., 366. 93-123. pomi, apple, 306. Internal breakdown, apple, 310. Internal cork (non-par.), apple, pear, 317. Lime sulfur injury, cherry, 315. 310. Little peach (virus), peach, 298. Interveinal chlorosis (non-par.),

chestnut, 338.

Lophodermium sp., Rhododendron spp., 372.

Lophodermium pinastri, Pinus spp., Melampsora bigelowii, Salix spp., 343. cassandrae, Picea spp., 348. 348. rhododendri, Rhododendron lini, flax, 228. catawbiense, 372. Linum lewisi, 228, 366. Lycogala epidendrum, Pinus spp., ribesii-purpureae, Salix sp., 343. 348. Salix bebbiana, 343. Salix scouleriana, 343. M Melampsorella cerastii, Abies lasiocarna, 346. Macrophoma candollei; Buxus semper-Cerastium arvense, 355. virens, 354. Melanopsichium austro-americanum, negundinis, Acer negundo, 337. Polygonum lapathifolium, 371. nervicola, Quercus spp., 342. Micrococcus sp., undetermined Gramineae, phlei, Phleum pratense, 246. Microorganisms and viruses, in Hawaii Macrophomina phaseoli, bean, 281. cowpea, 242. and their hosts, 38-82. honeydew melon, 265. Microsphaera sp., Euonymus spp., 360. Persian melon, 265. Symphoricarpos spp., 374. Strophostyles helvola, 241. Microsphaera alni, pecan, 324. Macrosporium sp., Dahlia sop., 357. Syringa vulgaris, 375. onion, 248. alni vaccinii, Catalpa spb., 338. diffusa, Symphoricarpos spp., 374. catalpae, Catalpa spp., 338. polonica, Hydrangea sp., 364. porri, onion, 248. sarcinula parasiticum, onion, 249. Mild mosaic (virus), potato, 292. Malformation, Italian prune, 313. Milo disease, reaction of sorghum Malnutrition (non-par.), cauliflower; varieties and hybrids to, 165-175. 256. Marasmius tritici, Agrostis alba, 244. Missing hills, hops, 328. Monilochaetes infuscans, sweetpotato, wheat, 231. 189, 272. Marginal chlorosis (non-par.), chest-Mosaic (virus), alfalfa, 240. nut, 338. Marssonia sp., crabapple, 336. bean, 283. green ash, 335. black raspberry, 322. hawthorn, 336. Boysenberry, 320. fraxini, see Piggotia fraxini. cabbage, 259. juglandis, Juglans spp., 340. Callicarpa americana, 354. Marssonina sp., Salix spp., 343. cantaloupe, 265. cauliflower, 256. kriegeriana, Salix exigua, 343. panattoniana, lettuce, 273. clover, 242. populi, Populus alba, 341. cowpea, 243. Populus tremuloides, 335, 336. cucumber, 267. Measles (non-par.), apple, 310. currant, 319. Melampsora spp., cottonwood, 335. Dahlia spo., 357. dry beans, 197. Populus spp., 336, 341. Populus balsamifera, 336. eggplant, 287. Populus candicans, 336. garden beet, 253. Populus carreiriana, 336. Gladiolus spp., 262. Melampsora abietis-caradensis, green beans, 196.

Populus sop., 341.

Melampsora aecidioides, Populus

alba, 341.

green peas, 198.

Lathyrus odoratus, 366.

Iris sp., 364.

Mosaic (virus), lettuce, 274. Lilium spp., 366. lima bean, 280. Melilotus alba, 240. Narcissus sp., 367. Ornithogalum aureum, 368. Paeonia spp., 369. pea, 285. peach, 298. Pelargonium spp., 370. pepper, 261. Petunia spp., 370. Phytolacca americana, 370. potato, 190. prune, 313. purple raspberry, 322. red raspberry, 322. Rosa spp., 373. Senecio cruentus, 374. soybean, 240. spinach, 294. squash, 268. strawberry, 304. sugar cane, 334. summer squash, 269. sweetpotato, 272. Tigridia sp., 375. tobacco, 202, 332. tomato, 193, 195, 279. turnip, 260. watermelon, 263. wheat, 233. winter squash, 269. Zinnia elegans, 377. Mottle leaf (virus), cherry, 315. Mount Shasta, California, host index to fungi of, 125-133. Nummularia discreta, apple, 307. Mucor spp., field corn, 236. Multiple bud (undet.), sugar cane, 334. Mycosphaerella sp., tomato, 276. brassicicola, Brassica oleracea var. broccoli, 254. Brussels sprouts, 259. cabbage, 257. cauliflower, 255. cercidicola, Cercis canadensis, 339. citrullina, watermelon, 262. winter squash, 269.

fragariae, strawberry, 208, 302.

Mycosphaerella pinodes, green peas, 198, 199. pea, 284. pomi, apple, 307. rosicola, Rosa spp., 373. rubi, blackberry, 319. Boysenberry, 320. raspberry, 321. youngberry, 322. rubina, raspberry, 320, 321. red raspbery, 321. sentina, pear, 317.

Naucoria cerealis, wheat, 231. Necrosis, tomato, 279. Nectria sp., Abies balsamea, 346. Abies concolor, 346. horsechestnut, 337. ornamentals, 349. Ulmus spp., 345. cinnabarina, currant, 319. Fagus spp., 339. Ligustrum amurense, 366. Ulmus spp., 345. ditissima, Acer spp., 337. rousselliana, Buxus semper virens, 354. Nematode conference, proceedings third national plant, 135-150. Nematodes, Citrus sinensis, 300. in Hawaii and their hosts, 76-82. Neofabraea malicorticis, apple, 307. perennans, apple, 307.

Nigrospora sphaerica, field corn, 235.

Net necrosis (virus), potato, 291.

Oidium sp., Hydrangea sp., 364. caricae, papaya, 300. Oil injury, apple, 311. Ophiobolus graminis, barley, 180. wheat, 231. miyabeanus, rice, 229. Ovularia asteris, Aster engelmanni, 352. obliqua, Rumex.sp., 373. Ovulinia azaleae, Azalea spp., 352.

Penicillium sp., Citrus spp., 300. cotton, 325, 326. cranberry, 322. field corn, 235, 236. prune, 313. Tulipa spo., 376. expansum, apple, 307. pear, 318. gladioli, Gladiolus spp., 362. Peronospora sp., Eriogonum blissianum, 360. Teucrium canadense, 375. antirrhini, Antirrhinum majus, 351. destructor, onion, 249. effusa, Chenopodium album, 355. spinach, 294. lamii, Lamium amplexicaule, 365. manshurica, soybean, 240. parasitica, cabbage, 257. Capsella bursa-pastoris, 355. cauliflower, 255. radish, 285. pisi, pea, 284. potentillae, Geum sp., 361. potentillae, Potentilla gracilis ssp. nuttallii, 371. schachtii, garden beet, 253. sugar beet, 333. tabacina, tobacco, 202, 330. trifoliorum, alfalfa, 238. Pestalotia sp., Rhodedendron spp.,372. guepini, Rhododendron spp., 372. Pezicula carpinea, Carpinus caroliniana, 338. Pezizella lythri, strawberry, 303. Phaeocryptopus gaumanni, Pseudotsuga taxifolia, 348. Phleospora aceris, Acer spp., 337. Acer saccharinum, 337. osmorrhizae, Osmorhiza divaricata, 363. Osmorhiza occidentalis, 368. Phloem necrosis (virus), Ulmus spp., 345. Phoma sp., peanut, 325. sugar beet, 333.

betae, sugar beet, 333.

Phoma destructiva, tomato, 276. fruit rot, tomato, 276. lathyrina, cowpes, 242. lingam, cabbage, 257. turnin, 259. Phoma persicae, peach, 296. terrestris, onion, 249. Phomopsis sp., Ilex spp., 340. callistephi, Callistephus chinensis, 354. gardeniae, Gardenia jasminoides, 361. juniperovora, cedar, 346. eastern red cedar, 335. Juniperus spp., 347. sojae, see Diaporthe sojae. stewartii, Cosmos spp., 357. vexans, eggplant, 286. Phony peach (virus), peach, 298. Phoradendron flavescens, Ulmus spp., 345. Phragmidium spp., Rosa spp., 373. americanum, Rosa setigera, 373. andersoni, Potentilla fruticosa, 371. ivesiae, Potentilla gracilis, 371. Potentilla gracilis ssp. nuttallii, Potentilla nuttallii, 371. montivagum, Rosa sp., 373. Rosa nutkana, 373. potentillae, Potentilla flabelliformis 371. Potentilla pensylvanica, 371. rosae-setigerae, Rosa spp., 373. Pestalozzia funerea, Thuja spp., 349. Phragmidium rubi-idaei, blackberry, 321. Rubus strigosus, 321. speciosum, Rosa spp., 373. Phyllachora trifolii, Trifolium hybridum, 241. Trifolium pratense, 241. Trifclium repens, 241. Phyllosticta spp., Acer.spp., 337. Amaranthus retroflexus, 350. Buxus sembervirens, -354. Caroinus caroliniana, 338. Catalpa sop., 338. crabapple, 336. Fraxinus spp., 340. Fraxinus velutina, 340. hawthorn, 336.

Phyllosticta sp., Hedera spp., 363. Phymatotrichum omnivorum, Populus spp. Phlox spp., 37C. 341. Sassafras albidum, 344. Pyracantha sp., 372. Ulmus spp., 345. Rhus spp., 343. antirrhini, Antirrhinum majus, 351. Rosa spp., 373. bumeliae, Bumelia lanuginosa, 338, Salix spp., 343. 353. Spiraea spp., 374. catalpae, Catalpa spp., 338. sweet clover, 240. celtidis, Celtis occidentalis, sweetpotato, 272. ·338. swiss chard, 254. gossypina, cotton, 326. Thuja orientalis, 349. hederaecola, Hedera helix, 363. tobacco, 331. ilicicola, Ilex spp., 340. tomato, 276. kalmicola, Kalmia latifolia, 365. Ulmus americana, 345. smilacis, Smilax glauca, 374. Zinnia elegans, 377. solitaria, apple, 204, 205, 307. Physalospora astragali, Astragalus straminella, rhubarb, 286. bisulcatus, 352. trollii, Trollius albiflorus, 376. Astragalus goniatus, 352. viridis, Fraxinus americana, 340. obtusa, apple, 204, 205, 308. Phymatotrichum omnivorum, alfalfa, Physarum spp., undetermined Gramineae, 239. 247. apricot, 312. cinereum, Cynodon dactylon, 244. Stenotaphrum secundatum, 246. Aster spp., 352. Physiological yellow leaf, cherry, 315. Baccharis neglecta, 353. Buxus sempervirens, 354. Physoderma menyanthis, Menyanthes trifoliata, 367. Cercis canadensis, 339. zeae-maydis, field corn, 236. Chrysanthemum spp., 356. Physopella fici, fig, 301. Cornus spp., 339. Phytomonas spp., Dianthus caryophyllus, cotton, 201, 327. cowpea; 242. 358. dry beans, 197. Elaeagnus argentea, 359. green beans, 196. Euonymus spp., 360. Syringa vulgaris, 375. fig; 301. Gelsemium sembervirens, 361. tobacco, 202. aceris, Acer spp., 337. Hibiscus syriacus, 363. aleuritidis, tung-oil tree, 325. Juglans rupestris, 340. andropogoni, sorghum, 230. Lantana camara, 365. Lantana sellowiana, 365. Sorghum vulgare sudanense, 246. angulata, tobacco, 331. Ligustrum sp., 366. atrofaciens, wheat, 231. Ligustrum amurense, 366. begoniae, Begonia spp., 353. Ligustrum lucidum compactum, berberidis, Berberis spp., 353. 366. campestris, Brassica oleracea var. Melia azedarach, 367. broccoli, 254. mulberry, 311. Brassica oleracea var. napookra, 271. brassica, 254. pear; 318. pecan; 324. cabbage, 258. peach 206, 296. cauliflower, 255. radish, 286. Philadelphus grandiflorus, turnip, 259. 370.

Phytomonas cerasi, Trifolium Phytomenas stewartii, popcorn, 237. hybridum, 241. sweet com, 188, 237. Trifolium pratense, 241. syringae, lima bean, 280. Trifolium repens, 241. plum, 316. coronafaciens, oats, 225. sweet cherry, 314. cucurbitae, squash, 268. Syringa vulgaris, 375. winter squash, 269. tabaci, tobacco, 331. delphinii, Delphinium spp., 358. tardicrescens, Iris sp., 364. translucens, barley, 227. fascians, Lathyrus odoratus, 365. flaccumfaciens, bean, 281, 282. translucens undulosa, wheat, 231. glycinea, soybean, 240. tumefaciens, Althaea rosea, 350. gummisudans, Gladiolus spp., 362. apple, 308. hederae, Hedera spp., 363. black raspberry, 321. insidiosa, alfalfa, 239. blackberry, 320. jaggeri, celery, 251. lachrymans, cucumber, 266. Boysenberry, 320. cotton, 32%. maculicolá, cabbage, 258. Euonymus fortunei var. radicans, cauliflower, 255. 360. malvacearum, cotton, 201, 327. grape, 323. hops, 328, 329. marginata, Gladiolus spp., 362. Ligustrum lucidum compactum, 366. medicaginis phaseolicola, bean, 282. peach, 296. pecan, 324. lima bean, 280. plum, 316. michiganensis, tomato, 193, 195, raspberry, 321. 276. mori, Morus alba tatarica, 311. Rosa spp., 373. mulberry, 335. Salix spp., 344. papavericola, Papaver spp., 369. vesicatoria, pepper, 260. tomato, 277. papulans, apple, 308. viburni, Viburnum spp., 377. pelargonii, Pelargonium spp., 369. phaseoli, bean, 282. viridilivida, lettuce, 273. lima bean, 280. woodsii, Dianthus caryophyllus, 358. pisi, green peas, 198. Phytophthora sp., Citrus grandis, 300. pea, 284. Callistephus chinensis, 354. pruni, apricot, 312. Lilium spp., 366. cherry, 314. rhubarb, 286. peach, 206, 296. strawberry, 208, 303. plum, 316. Syringa vulgaris, 375. cactorum, Antirrhinum majus, 351. rhizogenes, apple, 308. apple, 308. Rosa spp., 373. rubrilineans, sugar cane, 334. Cornus spp., 339. rubrisubalbricans, sugar cane, 334. Cornus florida, 357. Paeonia spp., 369. savastanoi, Nerium oleander, 368. sepedonica, potato, 191, 288. Rhododendron spp., 372. strawberry, 303. solanacearum, Dahlia spp., 357. potato, 191, 289. Syringa wilgaris, 375. tobacco, 331. cansici, pepper, 260. cryptogea richardiae, Zantedeschia tomato, 276. stewartii, corn, 186, 187. sp., 377. field corn, 236. erythroseptica, potato, 289.

Phytophthora infestans, potato, 190, Pratylenchus musicola, black walnut, 289. tomato, 277. megasperma, sweet clover, 240. parasitica, Citrus spp., 300. tomato, 277. parasitica nicotianae, tobacco, 331. phaseoli, lima bean, 280. Pierce's disease (virus), grapevines, Proceedings, third national plant Piggotia fraxini, ash, 336. Pink cherry (virus), saur cherry, Pink fruit, cherry, 315. Piricularia grisea, Digitaria sanguinalis, 245. oryzae, rice, 229. Pitya cupressi, Juniperus spp., 347. Plant hosts, microorganisms and viruses in Hawaii, 3-36. Plant pathogens and virus diseases, eastern Asia, host index, 93-123. Plasmodiophora brassicae, Brassica oleracea var. broccoli, cabbage, 258. cauliflower, 256. turnip, 259. Plasmopara geranii, Geranium carolinianum, 361. viticola, grape, 209, 323. Plen odomus destruens, sweetpotato, 272. Pleospora lycopersici, tomato, 277. Pod rot (undet.), peanut, 325. Podosphaera spp., peach, 296. leucotricha, apple, 308. oxyacanthae, cherry, 314. sour cherry, 314. western chokecherry, 342. Wickson plum, 316. Polyporus amarus, Libocedrus decurrens, 335. hirsutus, birch, 338. versicolor, Populus spp., 341. volvatus, Pinus contorta, 348. Poor growth, Tsuga spp., 349. Potash deficiency, cotton, 327. tobacco, 332.

146. boxwood, 146. fig, 146. olive, 146. pratensis, Buxus sempervirens, 354. Chrysanthemum spp., 356. cotton, 146, 147. Gilia rubra, 362. nematode conference, 135-150. Pseudoperonospora cubensis, cantaloupe, 264. cucumber, 266. squash, 268. watermelon, 263. humuli, hops, 328, 329. Pseudopeziza medicaginis, alfalfa, 239. ribis, currant, 319. gooseberry, 313. wild gooseberry, 318. trifolii, Trifolium pratense, 241. Psyllid yellows (psyllids), potato, 191, 292. tomato, 278. Puccinia spp., barley, 227. aemulans, Viguiera multiflora, 377. albulensis, Veronica wormskjoldii, 376. andropogonis, Andropogon furcatus, 244. Andropogon scoparius, 244. anomala, barley, 181, 227. antirrhini, Antirrhinum majus, 351. aristidae, Chenopodium rubrum, 355. Sarcobatus vermiculatus, 373. Triglochin maritima, 376. asparagi, asparagus, 252. atrofusca, Carex sp., 355. Carex deflexa, 355. balsamorhizae, Balsamorhiza sagittata, bartholomaei, Bouteloua curtipendula, 244. bistortae, Polygonum viviparum, 371. canaliculata, Cyperus esculentus, 357. caricis, gooseberry, 318. caricis grossulariata, Ribes aureum, 319. Ribes setosum, 319.

Puccinia chrysanthemi, Chrysanthemum spp., 356.

clintonii, Pedicularis paysoniana, 369.

convolvuli, Convolvulus arvensis, 356.

coronata, Calamagrostis rubescens, 244.

Elaeagnus canadensis, 359.

Koeleria cristata, 245.

oats, 185, 225.

Rhamnus cathartica, 343.

crandallii, Symphoricarpos albus, 375.

Symphoricarpos occidentalis, 375.

douglasii, Phlox glabrata, 370. extensicola, Carex festivella,

Erigeron salsuginosus, 359. extensicola asteris, Aster sp., 352.

Aster foliaceus var. apricus, 352.

Aster foliaceus var. canbyi, 352.

extensicola grossulariata,
Agoseris glauca, 350.

extensicola hieraciata, Hieracium scouleri, 363.

fraxinata, Fraxinus spn., 340. green ash, 335.

gayophyti, Gayophytum caesium, 361. glumarum, Elymus glaucus, 245. wheat, 231.

graminis, Agrostis alba, 244. barley, 181, 227.

oats, 184, 225.

rye, 183, 230.

wheat, 178, 179, 231.

graminis phlei-pratensis, Phleum pratense, 246.

grumosa, Zigadenus elegans, 378. harknessii, Stephanomeria pauciflora, 374.

helianthellae, Helianthella uniflora, 363.

helianthi, Helianthus spp., 363. sunflower, 238.

heterospora, Althaea rosea, 350.

Puccinia hieracii, Agoseris aurantiaca, 350.

Pyrrhopappus carolinianus, 372. Taraxacum paludosum var. vulgare, 375.

iridis, Iris sp., 364

jonesii, Lomatium sp., 366. leveillei, Geranium fremonti, 361.

malvacearum, Althaea rosea, 350.

Malva rotundifolia, 367.

menthae, mint, 330.

-Monarda mollis, 367.

monoica, Arabis sp., 352.

obliqua, Gonolobus laevis, 362. obtegens, Cirsium arvense, 356.

pimpinellae, Osmorhiza obtusa, 368.

plumbaria, Phlox multiflora, 370. podophylli, Podophyllum peltatum, 371.

purpurez, Sorghum halepense, 246. rubigo-vera, Acuilegia caerulea, 351.

Clematis ligusticifolia, 356.

Ranunculus cymbalaria, 372. Ranunculus glaberrimus, 372.

rubigo-vera agropyri, Thalictrum occidentale, 375.

rubigo-vera apocrypta, Hydrophyllum capitatum, 364.

rubigo-vera secalis, rye, 182, 230. rubigo-vera tritici, barley, 181, 227. wheat, 178, 179, 232.

smilacis, Smilax bona-nox, 374.

solheimii, Dodecatheon pauciflorum, 359.

sorghi, field corn, 236.

Sorghum vulgare sudanense, 246. sweet corn, 237.

suksdorfii, Agoseris aurantiaca, 350. tardissima, Arenaria congesta, 352.

virgata, Sorghastrum nutans, 246. windsoriae, Ptelea trifoliata, 342.

Triodia flava, 247.

xanthii, Ambrosia trifida, 350.

Xanthium sp., 377.

Pucciniastrum americanum, red raspberry, 321.

goeppertianum, Abies lasiocarpa, 346.

huckleberry, 304.

Vaccinium membranaceum, 376. myrtilli, Thuja canadensis, 349.

Pucciniastrum myrtilli, Vaccinium membranaceum, 376. pyrolae, Pyrola secunda, 372. Purple top (virus), potato, 190. Purple top wilt (undet.), potato, 293. Pyrenopeziza medicaginis, alfalfa, Pyrenophora teres, barley, 227. Pythium sp., Agrostis spp., 243. Antirrhinum majus, 351. corn, 187, 236. cucumber, 267. flax, 228. garden beet, 253. lettuce, 273. pea, 283. pepper, 260. Phaseolus aureus, 280. Pinus spp.; 348. rhubarb, 286. sugar beet, 333. sugar cane, 334. tobacco, 330. tomato, 277. watermelon, 263. wheat, 231. acanthicum, watermelon, 263. aphanidermatum, potato, 290. radish, 286. sweetpotato; 272. arrhenomanes, sorghum, 230. butleri, field corn, 236. debaryanum, dasheen, 263. oats, 1225. sugar beet, 333. debaryanum type, pepper, 261. irregulare, spinach, 294. root rot, milo, see milo disease. ultimum, cowpea, 242. lettuce, 273. pea, 284. spinach, 294. sweetpotato, 189, 272. R

Ramularia sp., Leersia hexandra, 245. armoraciae, horseradish, 235. asteris, Aster fromontii, 352. Ramularia betae, sugar beet, 333. celtidis, Celtis occidentalis, 338. · cynarae, globe artichoke, 270. · heraclei, Heracleum lanatum, 363. pastinacae, parsnip, 280. primulae, Primula sp., 371. smilacinae, Smilacina racemosa, 374. Ravenelia epiphylla, Tephrosia virginiana, 375. Razoumofskya sp., Pseudotsuga taxifolia, 348. Red stele, strawberry, 208. Red suture disease, peach, 298. Rehmiellopsis bohemica, Abies concolor, 346. Rhabdocline pseudotsugae, Pseudotsuga taxifolia, 348. Rhizoctonia sp., Azalea spp., 352. celery, 251. Chrysanthemum spp., 356. Dianthus caryophyllus, 358. Dolichos sphaerospermus, 359. Euphorbia spp., 360. garden beet, 253. Gardenia jasminoides, 361. Gladiolus spp., 362. pepper, 260. Pinus spp., 348. Stenotaphrum secundatum, 246. sugar beet, 333. tobacco, 330. undetermined Gramineae, 247. bataticola, Abutilon theophrasti, 350. cotton, 325, 327. cowpez, 242. Erigeron canadensis, 359. Lactuca serriola, 365. Lespedeza stipulacea, 238. lima bean, 280. Phaseolus aureus, 280. soybean, 240. sweetpotato, 272. see also Sclerotium bataticola. microsclerotia, bean, 282. oryzae, rice, 229. solani, cotton, 325, 327. Cynodon dactylon, 244.

Epidendrum sp., 359. garden beet, 253.

```
Rhizoctonia solani, Myrica spp., 367.
                                        pez, 284.
                                    Salt injury (non-par.), celery, 252.
     Papaver orientale, 369.
     Pelargonium spp., 369.
                                     Savoy disease (virus), sugar beet, 334.
     Petunia spp., 370.
                                     Scald (non-par.), apple, 310.
     potato, 190.
                                           onion, 250.
     rhubarb, 286.
                                      Scaly bark (virus ?), Citrus spp., 300.
     rice, 229.
                                      Schizophyllum spp., ornamentals, 349.
      sugar beet, 333.
                                     Sclerospora macrospora, oats, 225.
                                      Sclerotinia sp., Antirrhinum majus, 351.
      undetermined Gramineae, 247.
Rhizonus spp., cotton, 327.
                                            cherry, 314.
                                           peach, 206, 297.
     field corn, 235.
                                           Tulipa spp., 376.
     prune, 313.
      squash, 268.
                                         camelliae, Camellia japonica, 355.
      sweetpotato, 189, 272.
                                         carunculoides, Morus alba, 311.
   nigricans, peach, 296.
                                         convoluta, Iris sp., 364.
      strawberry, 303.
                                        fructicola, apple, 308.
      sweetpotato, 272.
                                            cherry, 207.
                                           peach, 206, 297.
   rot, peach, 206.
Rhynchosporium secalis, barley, 223.
                                           plum, 316.
      rye, 230.
                                           sour cherry, 314.
Rhytisma acerinum, Acer spp., 337.
                                           sweet cherry, 314.
   erythrosporum, Quercus
                                         gladioli, Gladiolus sp., 362.
         virginiana, 343.
                                         homeocarpa, undetermined Gramineae,
Ringspot (virus), Antirrhinum majus,
                                              . 247.
          351.
                                         laxa, almond, 324.
     apricot, 312.
                                           abricot, 312.
                                          cherry, 207.
      cabbage, 259.
     Dahlia spp., 357.
                                            neach, 206.
     tobacco, 332.
                                         sclerotiorum, bean, 282.
     tomato, 279.
                                            cabbage, 258.
Root diseases, strawberry, 208.
                                         carrot, 270.
Root girdle (undet.), carrot, 270.
                                         cauliflower, 256.
Rcot knot, peach, 206.
                                           cucumber, 267.
                                           Hulianthus app., 363.
     tobacco, 202.
Root rot, dry peans, 197.
                                           lettuce, 273.
                                         . Malvaviscus sp., 367.
     field corn, 236.
                                           Matthiola incana, 367.
     hops, 328.
    milo, see milo disease.
                                           Matthiola incana 'annua, 367.
     Primule sp., 371.
                                           Paeonia spo., 369.
     squash, 268.
                                           Dua, 285. .
      sugar beet, 200
                                           potato, 290.
Root and stem rots, green peas, 198,
                                         squash, 268.
                                            strawberry, 303.
           199.
Rosellinia necatrix, Osmanthus aqui-
                                            winter squash, 269.
          folium, 368.
                                         trifoliorum, alfalfa, 239.
Rosette (virus), peach, 299.
                                            clover, 241.
Rotylenchulus reniformis, cotton, 146. Sclerotium bataticola, corn, 187.
      root parasite, 143.
                                          field corn, 235.
Rugose mosaic (virus), potato, 292.
                                          sorghum, 230.
                                           see also Rhizoctonia bataticola.
Running-out, carnation varieties, 359.
```

Sclerotium cepivorum, onion, 249. delphinii, Ageratum sp., 350. Antirrhinum majus, 351. Aquilegia spp., 351. Campanula medium, 355. Delphinium spp., 358. Eupatorium sp., 360. Trillium sp., 376. Viola spp., 377. oryzae, see Leptosphaeria salvinii. rolfsii, Allium sativum, 250. bean, 282. carrot, 270. Delphinium spp., 358. eggolant, 286. garden beet, 253. Hydrangea sp., 364. Iris sp., 364. Lespedeza stipulacea, 238. Lippia canescens, 366. peanut, 325. pepper, 261. potato, 290. sugar beet, 333. sugar cane, 334. tomato, 277. Scolecotrichum graminis, Agrostis alba, 244. Phleum pratense, 246. Scorias spongiosa, Fagus spp., 339. Seed pièce decay, potato, 293. Seedling blights, cotton, 201, 327. rye, 183. Septobasidium sydowii, Celtis occidentalis, 338. Sentoria sp., crabapple, 336. Fraxinus nigra, 340. hawthorn, 336. Panicum clandestinum, 246. apii, celery, 251. apii-graveolentis, celery, 251. azaleae, Azalea spp., 353. chrysanthemella, Chrysanthemum spp., 356. cucurbitacearum, squash, 268. gladioli, Gladiclus spp., 362. helianthi, Helianthus spp., 363. lactucae, lettuce, 273. lactucicola, Lactuca canadensis, 365.

Septoria leaf spot, Chrysanthemum spo., lycopersici, tomato, 192, 194, 277. nodorum, wheat, 233. obesa, Chrysanthemum spp., 356. philadelphi, Philadelphus spp., 370. phlogis, Phlox drummondii, 370. pisi, pea, 285. ribis, gooseberry, 319. rumicis, Rumex venosus, 373. signalensis, Symphoricarpos oreophilus, 375. symphoricarpi, Symphoricarpos spp., 375. tritici, wheat, 233 vercnicae, Veronica spp., 376. Shoot rot, milo, see milo disease. Silvering of foliage, peach, 299. Skin breakdown, Florida oranges, 300. Sooty mold, hops, 329. Sorghum varieties and hybrids, reaction to milo disease, 165-175. Sorosporium ellisii, Andropogon spp., 244. reilianum, popcorn, 237. syntherismae, Cenchrus pauciflorus, Spanish measles (undet.), grape, 323. Sphaceloma fawcettii, Citrus spb., 300. menthae, mint, 329. symphoricarpi, Symphoricarpos spp., 375. viclae, Viola spp., 377. Sphacelotheca cruenta, Sorghum halepense, 246. sorghi, sorghum, 230. Sorghum halepense, 246. Sphaerella linorum, flax, 228. Sphaeropsis sp., Juniperus virginiana, 347. Nerium oleander, 368. Pinus spp., 348. Quercus spo., 343. ellisii, Pinus nigra austriaca, 348. Pinus sylvestris, 348. linearis, Carya sp., 338. Carya cordiformis, 338. ulmicola, Ulmus spp., 345. Sphaerotheca castagnei, Taraxacum spp., 375. humuli, black raspberry, 321.

Sphaerothe'ca humuli, blackberry, 320. dewberry, 320. Geranium fremonti, 361. hops, 329. red raspberry, 321. strawberry, 303. Viola son., 377. , humuli fuliginea, Plantago eriopoda, 371. mors-uvae, currant, 319. gooseberry, 319. pannosa, peach, 297. Rosa spp., 373. Spicaria spp., cotton, 327. Spindle tuber (virus), potato, 292. Spindling sprout, potato, 291. Split pit (non-par.), peach, 299. Spondylocladium atrovirens, potato, 291. Sporidesmium scorzonerae, salsify, Sporocybe azaleae, Azalea spp., 353. Spotted wilt (virus), celery, 252. Cosmos sulphureus, 357. lattuce, 274. pepper, 261. tomato, 195, 279. Zantedeschia sp., 377. Spotting of berries (undet.), grape, 323. Spray injury, apple, 311. cherry, 315. peach, 299. pear, 318. Staghead, Ulmus son., 345. Stalk rots, corn, 186, 187. field corn, 235. Stem cracking (non-par.), Euphorbia spo., 360. Stemphylium sarcinaeforme, clover, 241. solani; tomato, 278. Stereum purpureum, apple, 308. cherry, 314. rameale, birch, 338. Stippen, apple, see bitter pit. Stipple spot (undet.), peach, 299. Stony pit (virus), pear, 318. Storage rots, granefruit, 301.

Storage rots, sweetpotato, 189. Straighthead (non-par.), rice, 229. Streak (virus), black raspberry, 322. green peas, 198. Rosa spp., 373. Senecio cruentus, 374. tobacco, 332. tomato, 195, 279. Sulfur dioxide injury, grape, 323. Sulfur fruit scald, apple, 311. Sulfur 'leaf injury', apple, 311. Summaries of tobacco diseases in 1939, 330. Sunscald, apple, 311. asparagus, 253. Sunscorch, fig, 301. Synchytrium decipiens, Amphicarpa bracteata, 351. endobioticum, potato, 291.

Taphrina sp., sand cherry, 315. cerasi, cherry, 314. coerulescens, Quercus sp., 343. Quercus falcata f. triloba, 343. Quercus marilandica, 343. Quercus virginiana, 343. communis, blum, 316. crataegi, apple, 309. dearnessii, Acer spp., 337. deformans, peach, 206, 297. mirabilis, plum, 316. pruni, beach plum, 317. prune, 313. robinsoniana, Alnus incara, 337. Thecaphora deformans, Astragalus bisulcatus, 352. Thielaviopsis basicola, Lathyrus odoratiis, 366. tobanco, 202, 331. Tilletia spp., wheat, 178, 179. asperifolia, Muhlenbergia asperifolia, 367. horrida, rice, 229. levis, wheat, 233. tritici, wheat, 233. Tip blight (non-par.), Antirrhinum majus, 351.

Tipburn (non-par.), cabbage, 259.

Tipburn, cauliflower, 256.

lettuce, 273.

potato, 191, 292.

Tobacco mosaic, pepper, 261.

Tranzschelia pruni-spinosae, plum, 317.

prune, 313.

pruni-spinosae discolor, peach, 298.

Trochila populorum, Populus sop., 341.

Tubercularia fici, fig, 301.
Typhula sp., undetermined Gramineae,
247.

[]

Uncinula sp., Parthenocissus quinquefolia, 369.

necator, grape, 209, 323. salicis, Salix sp., 344.

Salix glaucophylla var., 344.

Urocystis cepulae, onion, 249. occulta, rye, 182, 230.

Uromyces betae, sugar beet, 333. caladii, Arisaema dracontium,

352.

caryophyllinus, Dianthus caryophyllus, 358.

coloradensis, vetch, 242.

elegans, Trifolium carolinianum, 241.

Trifolium procumbens, 241. graminicola, Panicum virgatum, 246.

Stillingia sylvatica, 374. hedysari-obscuri, Hedysarum marginatum, 363.

heterodermus, Erythronium grandiflorum, 360.

intricatus, Eriogonum campanulatum, 360.

jonesii, Ranunculus calthaeflorus, 372.

lannonicus, Oxytronis gracilis, 363.

lespedezae-procumbentis,

Lespedeza spp., 238.

minor, Trifolium parryi, 241. peckianus, Plantago aristata, 371.

Uromyces peckianus, Plantago purshii, 371.

phaseoli typica, bean, 283.

green beans, 196. lima bean, 280.

phaseoli vignae, compea, 243.

shearianus, Atriplex corrugata, 352.

spermacoces, Diodia teres, 359.

striatus, alfalfa, 239. trifolii, clover, 241.

trifolii fallens, clover, 241.

Trifolium pratense, 241.

trifolii hybridi, Trifolium hybridum, 242.

trifolii trifolii-repentis,

Trifolium repens, 242. zygadeni, Zigadenus gramineus, 378.

Ustilago spo., barley, 228.

Bromus spp., 244.

avenae, cats, 184, 226.

bromivora, Bromus spp., 244.

crusgalli, Echinochloa crusgalli, 245.

cynodontis, Cynodon dactylon, 245.

hordei, barley, 180, 228. rye, 230.

hypodytes, Agropyron spp., 243.

levis, oats, 184, 226.

lorentziana, Hordeum jubatum, 245.

nigra, barley, 228.

nuda, barley, 180, 228.

rabenhorstiana, Digitaria sanguin-

alis, 245.

striaeformis, Agrostis alba, 244.

Phleum pratense, 246.

tritici, wheat, 178, 179, 233.

zeae, corn, 186, 187.

field corn, 236. sweet corn, 188, 237.

V

Valsa spp., peach, 298.

Populus spp., 342.

celtidis, Celtis occidentalis, 338. cincta, peach, 298.

leucostoma, peach, 298.

Venturia sp., Pyracantha sp., 372. inaequalis, apple, 204, 205, 309.

Venturia inaequalis orbicularis, Crataegus spp., 339. byrina, pear, 203, 318. Vermicularia sp., Carya sop., 338. Ulmus spp., 345. Verticillium sp., Acer platanoides, 337. Acer pseudo-platanus, 337. Acer saccharum, 337. Antirrhinum majus, 351. Aucuba sp., 352. Berberis thunbergii, 353. blackberry, 320. Chrysanthemum morifolium, 356. eggpl.nt, 286. Phlox paniculata, 370. raspberry, 322. Rhus aromatica, 343. strawberry, 303. sugar boet, 334. tomate, 273. Ulmus spp., 345. Ulmus americana, 345. Viburnum tinus, 377. albo-atrum, Acer spo., 337. Antirrhinum majus, 351. apricot, 312. black raspberry, 321. Boysenberry, 320. Brassica oleracea var. navobrassica, 254. Chrysanthemum spo., 356. cotton, 327. Daldia spp., 357. eggolant, 296. okra, 271. Pasonia spo., 369. peach, 298. pepper, 261. Philak sop., 370. purple raspberry, 322. red raspberry, 322. Rhus spo., 343. sweetpotato, 272. tomato, 193, 195, 278. youngberry, 322. dahlise, Senecio cruentus, 374. Virus diseasos, boan, 283. cabbage, 259. celery, 252.

Virus discases, cherry, 315. cucumber, 267. dry beans, 197. garden beet, 253. green beans, 196. green peas, 198, 199. hops, 328, 329. lettuce, 273. peach, 206, 298. pepper, 261. potato, 190, 291. raspberry, 322 strawberry, 304. sugar beet, 334. tobacco, 332. tomato, 193, 195, 278. Virus diseases and plant pathogens, eastern Asia, host index, 93-123. Viruses, in Hawaii and their hosts, 82-84.

Wart (undet.), peach, 299. Wasting disease, Zostera marina, 378. Water core (non-par.), apple, 311. Weak hills, hops, 329. Weather injury, Abies spp., 346. apple, 311. cherry, 315. Delphinium spp., 358. Kalmia latifolia, 365. peach, 300. pear, 318. prune, 313. Rhododendron spp., 372. strawberry, 304. Taxus spp., 349. wheat, 233. Western celery mosaic (virus), celery, Western yellow mosaic (virus), celery, 252. White mottling of leaves (virus), Jatropha stimulosa, 365. White streak (virus), Narcissus sp., 367.

White tip (non-par.), rice, 229. Wind injury, apple, 311.

Wind injury, lettuce, 274. Wind and sand injury, tomato, 278. Winter injury, Abies soo., 346. alfalfa, 240. apple, 311. black raspberry, 322. grape, 323. Juniperus spp., 347. peach, 300. purple raspberry, 322. red raspberry, 322. strawberry, 304. sweet cherry, 316. Taxus spo., 349.

Witches' broom, alfalfa, 240. clover, 242. strawberry, 304. western yew, 349.

wheat, 233.

Χ

Xylaria mali, apple, 310.

Yellow dwarf (virus), potato, 292. Yellow mosaic (virus), bean, 283. Yellow-red virosis (virus), chokecherry, 299. peach, 299.

Prunus virginiana, 315, 342. Prunus virginiana var. demissa. 315.

Yellow spotting of leaves (undet.). peach, 299.

Yellow vein (undet.), peach, 299. Yellowing and stunting (non-par.) Delphinium spp., 358.

Yellows (undet.), Sassafras albidum, 344. Yellows (virus), Callistephus chinensis,

354. carrot, 270.

Cichorium endivia, 261.

Dianthus caryophyllus, 359.

peach, 299. strawberry, 304. tomato, 278.

Yellow top, tomato, 279.

Yellow dwarf (virus), onion, 250. Z-disease, potato, see Fusarium solani eumartii.

### ERRATA

On page 254 under Brassica campestris, read Erysiphe polygoni instead of Erysiphe graminis.

On page 278, 2d paragraph from bottom read psyllid instead of pyllid.

Low On page 283, last paragraph read Aphanomyces euteiches instead of Alphanomyces euteiches.

On page 298 last paragraph delete under phony peach "and in Oklahoma a trace loss." Phony peach was not reported from Oklahoma in 1939.

On page 340 under Fraxinus spp., read Cylindrosporium sp. instead of Clylindrosporium.

On page 351 under Antirrhinum majus, Sclerotinia sp., "S. delphinii" should be read Sclerotium delphinii, not Sclerotinia.

